arXiv:1105.4009v1 [math.AT] 20 May 2011

A DENSE SUBGROUP OF TOPOLOGICAL FUNDAMENTAL GROUPS OF QUOTIENT SPACES

HAMID. TORABI, ALI. PAKDAMAN AND BEHROOZ. MASHAYEKHY *

Department of Pure Mathematics, Center of Excellence in Analysis on Algebraic Structures, Ferdowsi University of Mashhad,
P.O.Box 1159-91775, Mashhad, Iran.

Abstract. In this paper, we show that the image of the topological fundamental group of a given space X is dense in the topological fundamental group of the quotient space X/A under the induced homomorphism of the quotient map, where A is a suitable subspace of X with some conditions on X. Also, we give some application to find out some properties for $\pi_1^{top}(X/A,*)$. In particular, we give some condition in which $\pi_1^{top}(X/A,*)$ is an indiscrete topological group.

1. Introduction and Motivation

Let X be a topological space and $\sim$ be an equivalent relation on X. Then one can consider the quotient topological space $X/\sim$ and the quotient map $p: X \longrightarrow X/\sim$. By applying the fundamental group functor on p there exists the induced homomorphism

$$p_*: \pi_1(X) \longrightarrow \pi_1(X/\sim).$$

It seems interesting to determine the image of the above homomorphism. Recently, Calcut, Gompf, and Mccarthy [5] proved that if X is a locally path connected topological space partitioned into connected subsets and if the associated quotient space $X/\sim$ is semi-locally simply connected, then the induced homomorphism, p_*, of fundamental groups is surjective for each choice of base point $x \in X$. They inspired the problem by one of the Arnold's problems on orbit spaces of vector fields on manifolds [5, Section 3.6].

D. Biss [2] introduced the topological fundamental group $\pi_1^{top}(X,x)$ of a based space (X,x) as the fundamental group $\pi_1(X,x)$ with the quotient topology of the loop space $\Omega(X,x)$ with respect to the canonical function $\pi: \Omega(X,x) \longrightarrow \pi_1(X,x)$ identifying path components. It is known that this construction gives rise a homotopy invariant functor $\pi_1^{top}: htop_* \longrightarrow QTG$ from the homotopy category of based spaces to the category of quasitopological groups and continuous homomorphism [3]. By applying the above functor on the quotient map $p: X \longrightarrow X/\sim$, we have a continuous homomorphism $p_*: \pi_1^{top}(X) \longrightarrow \pi_1^{top}(X/\sim)$.

Let X be a topological space and $A_1, A_2, ..., A_n$ be a finite collection of its subsets. The quotient space $X/(A_1, ..., A_n)$ is obtained from X by identifying each of the sets A_i to a point. Now, let (A,a) be a pointed subspace of (X,a) and $p: (X,a) \longrightarrow (X/A,*)$ be the associated quotient map. In this paper, first we prove that if A is an open subset of X such that the closure of A, $\overline{A}$, is path connected, then the image of p_* is dense in $\pi_1^{top}(X/A,*)$ and by this we show that the image of p_* is dense in $\pi_1^{top}(X/(A_1,A_2,...,A_n),*)$, where the A_i's are open subsets of X with path connected closure and $p: X \longrightarrow X/(A_1,A_2,...,A_n)$ is the associated quotient map. Second, we prove that if A is a closed subset of a locally path connected, first countable space X, then the image of p_* is also dense in $\pi_1^{top}(X/A,*)$. By two previous results we can show that the image

---

2010 *Mathematics Subject Classification.* 54B15; 55P65; 55Q52; 55Q70.

Key words and phrases. Topological fundamental group, Quotient map, Dense subgroup.

*Corresponding author.

E-mail addresses: hamid_torabi86@yahoo.com; alipaky@yahoo.com and bmashf@um.ac.ir.

of p_* is dense in $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$, where X is first countable, connected, locally path connected and A_i's are open or closed subsets of X with disjoint path connected closure. Moreover, we give some conditions in which p_* is an epimorphism. Also, by some examples, we show that p_* is not necessarily onto. Finally, we give some applications of the above results to find out some properties of the topological fundamental group of the quotient space $X/(A_1, A_2, ..., A_n)$. In particular, we prove that with the recent assumptions on X and the A_i's, $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is an indiscrete topological group when X is simply connected.

2. NOTATIONS AND PRELIMINARIES

For a pointed topological space (X, x), by a path we mean a continuous map $\alpha : [0, 1] \longrightarrow X$. The points $\alpha(0)$ and $\alpha(1)$ are called the initial point and the terminal point of α, respectively. A loop α is a path with $\alpha(0) = \alpha(1)$. For a path $\alpha : [0, 1] \longrightarrow X$, α^{-1} denotes a path such that $\alpha^{-1}(t) = \alpha(1 - t)$, for all $t \in [0, 1]$. Denote $[0, 1]$ by I, two paths $\alpha, \beta : I \longrightarrow X$ with the same initial and terminal points are called homotopic relative to end points if there exists a continuous map $F : I \times I \longrightarrow X$ such that

$$F(t, s) = \begin{cases} \alpha(t) & s = 0 \\ \beta(t) & s = 1 \\ \alpha(0) = \beta(0) & t = 0 \\ \alpha(1) = \beta(1) & t = 1. \end{cases}$$

The homotopy class containing a path α is denoted by $[\alpha]$. Since most of the homotopies that appear in this paper have this property and end points are the same, we drop the term "relative homotopy" for simplicity. For paths $\alpha, \beta : I \longrightarrow X$ with $\alpha(1) = \beta(0)$, $\alpha * \beta$ denotes the concatenation of α and β, that is, a path from I to X such that $(\alpha * \beta)(t) = \alpha(2t)$, for $0 \leq t \leq 1/2$ and $(\alpha * \beta)(t) = \beta(2t - 1)$, for $1/2 \leq t \leq 1$.

For a space (X, x), let $\Omega(X, x)$ be the space of based maps from I to X with the compact-open topology. A subbase for this topology consists of neighborhoods of the form $\langle K, U\rangle = \{\gamma \in \Omega(X, x) \mid \gamma(K) \subseteq U\}$, where $K \subseteq I$ is compact and U is open in X. When X is path connected and the basepoint is clear, we just write $\Omega(X)$ and we will consistently denote the constant path at x by e_x. The topological fundamental group of a pointed space (X, x) may be described as the usual fundamental group $\pi_1(X, x)$ with the quotient topology with respect to the canonical map $\Omega(X, x) \longrightarrow \pi_1(X, x)$ identifying homotopy classes of loops, denoted by $\pi_1^{top}(X, x)$. A basic account of topological fundamental groups may be found in [2], [4] and [3]. For undefined notation, see [11].

Definition 2.1. *([1]) A quasitopological group G is a group with a topology such that inversion $G \longrightarrow G$, $g \longrightarrow g^{-1}$, is continuous and multiplication $G \times G \longrightarrow G$ is continuous in each variable. A morphism of quasitopological groups is a continuous homomorphism.*

Theorem 2.2. *([4]) π_1^{top} is a functor from the homotopy category of based topological spaces to the category of quasitopological group.*

A space X is *semi-locally simply connected* if for each point $x \in X$, there is an open neighborhood U of x such that the inclusion $i : U \longrightarrow X$ induces the zero map $i_* : \pi_1(U, x) \longrightarrow \pi_1(X, x)$ or equivalently a loop in U can be contracted inside X.

Theorem 2.3. *([4]) Let X be a path connected space. If $\pi_1^{top}(X, x)$ is discrete for some $x \in X$, then X is semi-locally simply connected. If X is locally path connected and semi-locally simply connected, then $\pi_1^{top}(X, x)$ is discrete for all $x \in X$.*

3. MAIN RESULTS

In this section, (A, a) is a pointed subspace of (X, a), $p : (X, a) \longrightarrow (X/A, *)$ is the canonical quotient map so that $q := p|_{X-A} : X - A \longrightarrow X/A - \{*\}$ is a homeomorphism. Also, by applying the functor π_1^{top} on p we have a continuous homomorphism $p_* : \pi_1^{top}(X, a) \longrightarrow \pi_1^{top}(X/A, *)$.

Lemma 3.1. *If A is an open subset of X, then any loops $\alpha : I \longrightarrow \overline{\{*\}} \subseteq X/A$ based at $*$ is nullhomotopic.*

Proof. Define $F : I \times I \longrightarrow X/A$ by

$$F(t,s) = \begin{cases} \alpha(t) & s = 0 \\ * & s > 0. \end{cases}$$

If we prove that F is continuous, then F is a homotopy between α and e_*. For this let U be an open set in X/A, we show that $F^{-1}(U)$ is open in $I \times I$.
Case 1: If $* \in U$, then $F^{-1}(U) = F^{-1}(\{*\}) \cup (\alpha^{-1}(U) \times \{0\})$

$$= ((I \times (0,1]) \cup (\alpha^{-1}(U) \times \{0\})$$

$$= (I \times (0,1]) \cup (\alpha^{-1}(U) \times I)$$

which is open in $I \times I$.
Case 2: If $* \notin U$, then $U \cap \partial\overline{\{*\}} = \varnothing$ since if there exists $x \in \partial\{*\}$ such that $x \in U$, then $\{*\} \cap U \neq \varnothing$ which is a contradiction. Since $U \cap \overline{\{*\}} = (U \cap \{*\}) \cup (U \cap \partial\{*\}) = \varnothing$ and $\alpha(I) \subseteq \overline{\{*\}}$, hence $F^{-1}(U) = \varnothing$. ☐

Theorem 3.2. *Let A be an open subset of X such that $\overline{A}$ is path connected, then for each $a \in A$ the image of p_* is dense in $\pi_1^{top}(X/A,*)$ i.e:*

$$\overline{p_*\pi_1^{top}(X,a)} = \pi_1^{top}(X/A,*).$$

Proof. Step One: First we show that for every loop $\alpha : (I, \partial I) \longrightarrow (X/A, *)$ such that $\alpha^{-1}(\{*\}^c)$ is connected, $[\alpha] \in Im(p_*)$. By assumption and openness of $\{*\}$ in X/A, there exists $s_1, s_2 \in (0,1)$ such that $\alpha^{-1}(\{*\}^c) = [s_1, s_2]$. Since $\alpha^{-1}(\{*\}^c)$ is compact subset of I, it suffices to let $s_1 = inf\{\alpha^{-1}(\{*\}^c)\}$ and $s_2 = sup\{\alpha^{-1}(\{*\}^c)\}$. Let $\overline{\alpha} : [s_1, s_2] \longrightarrow X$ by $\overline{\alpha}(t) = q^{-1}(\alpha(t))$, then $\overline{\alpha}(s_1), \overline{\alpha}(s_2) \in \overline{A}$. Since if G is an open neighborhood of $\overline{\alpha}(s_1)$ and $G \cap A = \varnothing$, then $q(G) = p(G)$ is an open neighborhood of $\alpha(s_1)$. Using continuity of α, there exists an open neighborhood J of s_1 in I such that $\alpha(J) \subseteq G$. On the other hand, by definition of s_1, for all $s < s_1$, $\alpha(s) = *$ which implies that $* \in q(G)$ which is a contradiction since $* \notin Im(q)$. Similarly $\alpha(s_2) \in \overline{A}$. Since $\overline{A}$ is path connected, there exist two paths $\lambda_1 : [0, s_1] \longrightarrow \overline{A}$ and $\lambda_2 : [s_2, 1] \longrightarrow \overline{A}$ such that $\lambda_1(0) = \lambda_2(1) = a$, $\lambda_1(s_1) = \overline{\alpha}(s_1)$ and $\lambda_2(0) = \overline{\alpha}(s_2)$. Define $\widetilde{\alpha} : I \longrightarrow X$ by

$$\widetilde{\alpha}(t) = \begin{cases} \lambda_1(t) & 0 \leq t \leq s_1 \\ \overline{\alpha}(t) & s_1 \leq t \leq s_2 \\ \lambda_2(t) & s_2 \leq t \leq 1. \end{cases}$$

By gluing lemma $\widetilde{\alpha}$ is continuous, so it remains to show that $p \circ \widetilde{\alpha} \simeq \alpha \;\; rel\{*\}$. Put $\alpha_1 = \alpha|_{[0,s_1]}$, $\alpha_2 = \alpha|_{[s_2,1]}$ and let $\varphi_1 : [0,1] \longrightarrow [0, s_1]$ and $\varphi_2 : [0,1] \longrightarrow [s_2, 1]$ be linear homeomorphisms such that $\varphi_1(0) = 0$ and $\varphi_2(0) = s_2$, then $p \circ \lambda_i \circ \varphi_i \simeq \alpha_i \circ \varphi_i,\; rel\{0,1\}$ since the $(p \circ \lambda_i \circ \varphi_i) \circ (\alpha_i \circ \varphi_i)^{-1}$'s are loops in $\overline{\{*\}}$ which by Lemma 3.1 are nullhomotopic.
Step Two: By continuity of α, $\alpha^{-1}(\{*\}^c)$ is a closed subset of I. Connected subsets of I are intervals or one point sets, also connected components of $\alpha^{-1}(\{*\}^c)$ are closed in $\alpha^{-1}(\{*\}^c)$ and so compact in I. Therefore a component of $\alpha^{-1}(\{*\}^c)$ is either closed interval or singleton. Given $[\alpha] \in \pi_1(X/A,*)$, we show that there exists a sequence of homotopy classes of loops $\{[\alpha_n]\}_{n \in N}$ in $Im(p_*)$ such that $[\alpha_n] \longrightarrow [\alpha]$ in $\pi_1^{top}(X/A,*)$.

We claim that the number of non-singleton components of $\alpha^{-1}(\{*\}^c)$ is countable. Let S be the union of singleton components of $\alpha^{-1}(\{*\}^c)$ and for each $n \in \mathbb{N}$, B_n be the set of non-singleton components of $\alpha^{-1}(\{*\}^c)$ with length at least $1/n$. Each B_n is finite since if B_n is infinite, then it has at least $n+1$ members. Therefore $\bigcup_{C \in B_n} C \subseteq \alpha^{-1}(\{*\}^c) \subseteq I$ which implies that

$$(n+1) \times 1/n \leq \sum_{C \in B_n} diam(C) \leq diam(I) = 1$$

which is a contradiction. Thus each B_n is finite which implies that $B = \bigcup_{n \in \mathbb{N}} B_n$ is countable. Rename elements of B by $I_i = [a_i, b_i]$, $i \in J = \{1, 2, ..., s\}$, where $s = |B|$ if B is finite and $i \in \mathbb{N} = J$ if B is infinite. For every

$n \in J$ define

$$\alpha_n(t) = \begin{cases} \alpha(t) & t \in \bigcup_{i=1}^{n}[a_i, b_i] \\ * & otherwise. \end{cases}$$

If B is finite, put $\alpha_n = \alpha_s$, for every $n > s$. We claim that the α_n's are continuous. For, if $V \subseteq X/A$ is open, then

i) If $* \in V$, then

$$\alpha_1^{-1}(V) = [0, a_1) \cup (b_1, 1] \cup \alpha|_{[a_1,b_1]}^{-1}(V)$$

which by continuity of α is open in I.

ii) If $* \notin V$, then we show that $\alpha_1^{-1}(V) = \alpha|_{[a_1,b_1]}^{-1}(V) = \alpha|_{(a_1,b_1)}^{-1}(V)$ which guaranties $\alpha^{-1}(V)$ is open. For this it suffices to show that $\alpha_1(a_1), \alpha_1(b_1) \notin V$.

For each $n \in \mathbb{N}$, $\alpha(a_n), \alpha(b_n) \in \overline{\{*\}}$ and $\{\alpha(a)|\ a \in S\} \subseteq \overline{\{*\}}$. For, if G is an open neighborhood of $\alpha(a_n)$, then $\alpha^{-1}(G)$ is an open neighborhood of a_n, so there exists $\varepsilon > 0$ such that $(a_n - \varepsilon, a_n + \varepsilon) \subseteq \alpha^{-1}(G)$ or equivalently $\alpha((a_n - \varepsilon, a_n + \varepsilon)) \subseteq G$. If $* \notin G$, then $(a_n - \varepsilon, a_n + \varepsilon) \subseteq \alpha^{-1}(\{*\}^c)$ which is a contradiction since $[a_n, b_n]$ is a connected component of $\alpha^{-1}(\{*\}^c)$. Similarly $\alpha(b_n) \in \overline{\{*\}}$ for each $n \in \mathbb{N}$ and $\alpha(S) \subseteq \overline{\{*\}}$. Thus if $\alpha_1(b_1) = \alpha(b_1) \in V$, then V must meet $\{*\}$ which is a contradiction since $* \notin V$. Therefore α_1 is a continuous loop such that $[\alpha_1] \in Im(p_*)$. Similarly all the α_n's are continuous. Also, for every $n \in \mathbb{N}$, $[\alpha_n]$ is a product of n homotopy classes of loops which are similar to loops introduced in Step 1. This implies that $[\alpha_n] \in Im(p_*)$ since $Im(p_*)$ is a subgroup.

Now we show that the sequence $\{\alpha_n\}$ converges to α. Let $\alpha \in \langle K, U\rangle$, where K is a compact subset of I and U is an open subset of X/A, then

i) If $* \in U$, then for each $n \in \mathbb{N}$, $\alpha_n \in < K, U >$ since for each $t \in K$, $\alpha_n(t)$ is $\alpha(t)$ or $*$ which in both cases $\alpha(t) \in U$.

ii) If $K \subseteq \alpha^{-1}(\{*\}^c)$ and $K \cap S \neq \varnothing$, then there exists $a \in K \cap S \subseteq \alpha^{-1}(U)$, so $\alpha(a) \in U$, but $\alpha(a) \in \overline{\{*\}}$ and U is open. Thus $* \in U$ and by (i), for each n, $\alpha_n(K) \subseteq U$.

iii) If $K \subseteq \bigcup_{n=1}^{\infty} I_n$ and there exist $n_1, n_2, ..., n_s$ such that $K \subseteq \bigcup_{i=1}^{s} I_{n_i}$, then by definition of the α_n's, for each $n \geq max\{n_1, ..., n_s\}$ we have $\alpha_n(K) \subseteq U$.

vi) If $K \subseteq \bigcup_{n=1}^{\infty} I_n$ and there exists an infinite subsequence $\{I_{n_r}\}$ such that $K \cap I_{n_r} \neq \varnothing$, then there is a sequence $\{x_{n_r} | x_{n_r} \in K \cap I_{n_r}\}$ such that it has a subsequence $\{x_{n_{r_s}}\}$ converges to an element of K, b say, by compactness of K. Since $b \in K \subseteq \alpha^{-1}(U)$, there exists $\varepsilon > 0$ such that $(b - \varepsilon, b + \varepsilon) \subseteq \alpha^{-1}(U)$. Also, there exists s_0 such that for each $s \geq s_0$, $x_{n_{r_s}} \in (b - \varepsilon/2, b + \varepsilon/2)$ since $x_{n_{r_s}} \longrightarrow b$. Since $diam(I_n) \longrightarrow 0$, there is n_0 such that for each $n \geq n_0$, $diam(I_n) < \varepsilon/2$. Choose $s_1 \in N$ such that $s_1 \geq s_0$ and $n_{r_{s_1}} \geq n_0$, then $x_{n_{r_{s_1}}} \in (b - \varepsilon/2, b + \varepsilon/2)$. Also $x_{n_{r_{s_1}}} \in K \cap I_{n_{r_{s_1}}}$ and $diam(I_{n_{r_{s_1}}}) < \varepsilon/2$, so $a_{n_{r_{s_1}}} \in I_{n_{r_{s_1}}} \subseteq (b - \varepsilon, b + \varepsilon) \subseteq \alpha^{-1}(U)$ which implies that $\alpha(a_{n_{r_{s_1}}}) \in U$ and therefore $* \in U$ since $\alpha(a_{n_{r_{s_1}}}) \in \partial(\{*\})$. Using (i) $\alpha_n(K) \subseteq U$, for each $n \in \mathbb{N}$.

□

Definition 3.3. *Let X be a topological space and $A_1, A_2, ..., A_n$ be any subsets of X, $n \in \mathbb{N}$. By the quotient space $X/(A_1, ..., A_n)$ we mean the quotient space obtained from X by identifying each of the sets A_i to a point. Also, we denote the associated quotient map by $p : X \longrightarrow X/(A_1, A_2, ..., A_n)$.*

Corollary 3.4. *If A_1, A_2 are open subsets of a path connected space X such that $\overline{A_1}, \overline{A_2}$ are path connected. Then for every $a \in A_1 \cup A_2$,*

$$\overline{p_*\pi_1^{top}(X, a)} = \pi_1^{top}(X/(A_1, A_2), *).$$

Proof. We can assume that the A_i's are disjoint. If they are not disjoint, the result follows from Theorem 3.2. Let $p_1 : X \longrightarrow X/A_1$, $p_2 : X/A_1 \longrightarrow X/(A_1, A_2)$ be associated quotient maps and $a_1 = a \in A_1$. By Theorem 3.2, $\overline{(p_1)_*\pi_1^{top}(X, a_1)} = \pi_1^{top}(X/A_1, *_1)$, where $*_1 = p_1(a_1)$. Since X is path connected, so is X/A_1. Also, $p_1(A_2)$ is an open subset of X/A_1 and the closure of $p_1(A_2)$ in X/A_1 is path connected. Let $a_2 \in p_1(A_2)$, then $\overline{(p_2)_*\pi_1^{top}(X/A_1, a_2)} = \pi_1^{top}(X/(A_1, A_2), *_2)$, where $*_2 = p_2(a_2)$. Since X/A_1 is path connected, there exists a homeomorphism $\varphi_1 : \pi_1^{top}(X/A_1, *_1) \longrightarrow \pi_1^{top}(X/A_1, a_2)$ by $\varphi_1([\alpha]) = [\gamma * \alpha * \gamma^{-1}]$, where γ is a path from a_2

to $*_1$. We have $\overline{(p_2)_* \circ \varphi_1 \circ (p_1)_*(\pi_1^{top}(X,a))} \supseteq ((p_2)_* \circ \varphi_1)(\overline{(p_1)_*(\pi_1^{top}(X,a))}) = ((p_2)_* \circ \varphi_1)(\pi_1^{top}(X/A_1,*_1)) = (p_2)_*\pi_1^{top}(X/A_1,a_2) = Im(p_2)_*$ which implies that $Im(p_2)_* \circ \varphi_1 \circ (p_1)_*$ is dense in $\pi_1(X/(A_1,A_2),*_2)$. If $\gamma' = p_2 \circ \gamma$, then $\varphi_2 : \pi_1^{top}(X/(A_1,A_2),*_2) \longrightarrow \pi_1^{top}(X/(A_1,A_2),*_1)$ by $\varphi_2([\alpha]) = [\gamma'^{-1} * \alpha * \gamma']$ is a homeomorphism. Hence $Im(\varphi_2 \circ (p_2)_* \circ \varphi_1 \circ (p_1)_*)$ is dense in $\pi_1^{top}(X/(A_1,A_2),*_1)$. Moreover $\varphi_2 \circ (p_2)_* \circ \varphi_1 \circ (p_1)_* = p_*$ which implies that $Im(p_*)$ is dense, as desired. □

By induction and Corollary 3.4, we have the following results.

Corollary 3.5. *Let $A_1, A_2, ..., A_n$ be open subsets of a path connected space X such that the $\overline{A_i}$'s are path connected for every $i = 1, 2, ..., n$. Then for every $a \in \bigcup_{i=1}^n A_i$,*

$$\overline{p_*\pi_1^{top}(X,a)} = \pi_1^{top}(X/(A_1,A_2,...,A_n),*).$$

Corollary 3.6. *Let $A_1, A_2, ..., A_n$ be open subsets of a connected, locally path connected space X such that the $\overline{A_i}$'s are path connected for every $i = 1, 2, ..., n$. If $X/(A_1,A_2,...,A_n)$ is semi-locally simply connected, then for each $a \in \bigcup_{i=1}^n A_i$, $p_* : \pi_1^{top}(X,a) \longrightarrow \pi_1^{top}(X/(A_1,A_2,...,A_n,*)$ is an epimorphism.*

Proof. Let U be an open neighborhood of $\bar{x} \in (X/A_1) \setminus \overline{\{*\}}$. Since X is locally path connected, there is a path connected open neighborhood $\widetilde{U} \subseteq p^{-1}(U)$ of $x = q^{-1}(\bar{x})$ such that $\widetilde{U} \cap A_1 = \varnothing$. Then $V := p(\widetilde{U}) = q(\widetilde{U}) \subseteq U$ is a path connected open neighborhood of $\bar{x}$. X/A_1 is locally path connected at $*$ since $\{*\}$ is an open subset of X/A_1. Let U be an open neighborhood of $\bar{x} \in \partial(\{*\})$, then there exists a path connected open neighborhood $\widetilde{U} \subseteq p^{-1}(U)$ of x. Since $p^{-1}(p(\widetilde{U})) = \widetilde{U} \cup A_1$, $p(\widetilde{U})$ is a path connected open neighborhood of $\bar{x}$ in U. Therefore X/A_1 is locally path connected. Similarly $X/(A_1,A_2,...,A_n)$ is connected, locally path connected. Since $X/(A_1,A_2,...,A_n)$ is a connected, semi-locally simply connected and locally path connected space, by Theorem 2.3, $\pi_1^{top}(X/(A_1,A_2,...,A_n),*)$ is a discrete topological group which implies that $Im(p_*) = \pi_1^{top}(X/(A_1,A_2,...,A_n),*)$ by Corollary 3.5. □

In the following example we show that with assumption of Theorem 3.2, p_* is not necessarily onto.

Example 3.7. *Let $A_n = \{1/(2n-1), 1/2n\} \times [0, 1+1/2n] \bigcup [1/2n, 1/2n-1] \times \{1+1/2n\}$ for each $n \in \mathbb{N}$. Consider $X = (\bigcup_{n\in\mathbb{N}} A_n) \bigcup \{0\} \times [0,1] \bigcup [0,1] \times \{0\}$ with $a = (0,0)$ as the base point and $A = \{(x,y) \in X \mid y < 1\}$ (see Figure 1). A is an open subset of X with path connected closure. Assume $I_n = (1/2+1/2(n+1), 1/2+1/2n]$ and f_n be a homeomorphism from I_n to $A_n - \{(1/2n, 0)\}$ for every $n \in \mathbb{N}$. Define $f : I \longrightarrow X$ by*

$$f(t) = \begin{cases} \text{the point } (0,2t) & t \in [0,1/2], \\ f_n(t) & t \in I_n. \end{cases}$$

We claim that $\alpha = p \circ f$ is a loop in X/A at $*$. It suffices to show that α is continuous on $t = 1/2$ and boundary points of I_n's since f is continuous on $[0,1/2)$ and by gluing lemma on $\bigcup int(I_n)$. Since α is locally constant at $t = 1/2 + 1/2n$ for each $n \in \mathbb{N}$, α is continuous at boundary points of I_n. For each open neighborhood G of $f(1/2) = (0,1)$ in X, there exists $n_0 \in \mathbb{N}$ such that G contains $A_n \bigcap A^c$ for $n > n_0$. Therefore continuity at $t = 1/2$ follows from $\alpha(1/2) \in \overline{\{*\}}$. Now let $B \subseteq \mathbb{N}$ and define

$$g_B(t) = \begin{cases} (p \circ f)(t) & t \in \bigcup_{m\in B} I_m, \\ * & \text{otherwise}. \end{cases}$$

Then g_B is continuous and for $B_1, B_2 \subseteq \mathbb{N}$ such that $B_1 \neq B_2$, $[g_{B_1}] \neq [g_{B_2}]$ which implies that $\pi_1(X/A,*)$ is uncountable. But by compactness of I, a given path in X can traverse finitely many of the A_n's and therefore $\pi_1(X,a)$ is a free group on countably many generators which implies that p does not induce a surjection of fundamental groups.

Let (X,x) be a pointed topological space such that $\{x\}$ is closed. If $\alpha : [0,1] \longrightarrow X$ is a loop in X based at x, then $\alpha^{-1}(\{x\})$ is a closed subset of $[0,1]$. Its complement, $\alpha^{-1}(\{x\}^c)$ is therefore the union of a countable collection of disjoint open intervals. We denote this collection of intervals by W_α.



FIGURE 1.

Definition 3.8. *Let (X, x) be a pointed topological space. A loop α in X based at x is called* **semi-simple** *if $W_\alpha = \{(0,1)\}$ and is called* **geometrically simple** *if W_α has one element. If W_α is finite, then the loop α is called* **geometrically finite** [10].

Lemma 3.9. *Every geometrically simple loop is homotopic to a semi-simple loop.*

Proof. Let α be a geometrically simple loop at $x \in X$. Then there are $r, s \in [0,1]$ such that $\alpha^{-1}(\{x\}^c) = (r,s)$ and $\alpha(r) = \alpha(s) = x$. Let $\beta := \alpha|_{[r,s]}$ and $\varphi : [0,1] \longrightarrow [r,s]$ be a linear homeomorphism, then $\beta \circ \varphi$ is a semi-simple loop at x and $\alpha \simeq \beta \circ \varphi$. $\square$

In the sequel, for a semi-simple loop $\alpha : I \longrightarrow X/A$ denote $\widetilde{\alpha} = q^{-1} \circ \alpha|_{(0,1)} : (0,1) \longrightarrow (X - A)$, where A is a closed subset of a topological space X.

Lemma 3.10. *Let $A \subseteq X$ be a closed subset of X and α be a semi-simple loop at $*$ in X/A. If $lim_{t\to 0}\widetilde{\alpha}(t)$ and $lim_{t\to 1}\widetilde{\alpha}(t)$ do not exist, then for each $t_0 \in (0,1)$, there are $b_0, b_1 \in A$ such that b_0 is a limit point of $\widetilde{\alpha}((0,t_0))$ and b_1 is a limit point of $\widetilde{\alpha}((t_0,1))$.*

Proof. Let $t_0 \in (0,1)$ and by contradiction suppose each $b \in A$ has an open neighborhood G_b such that $G_b \cap \widetilde{\alpha}((0,t_0)) = \varnothing$, then $G = \bigcup_{b\in A} G_b$ is an open neighborhood of A and so $p(G)$ is an open neighborhood of $*$ (since $p^{-1}(p(G)) = G$) such that does not intersect $\alpha((0,t_0))$ which is a contradiction to continuity of α. $\square$

Theorem 3.11. *If A is a closed path connected subset of a locally path connected space X such that every point of A has a countable local base in X, then for each $a \in A$*

$$\overline{p_*\pi_1^{top}(X,a)} = \pi_1^{top}(X/A, *).$$

Proof. Step One: Let $[\alpha] \in \pi_1(X/A, *)$, where α is a semi-simple loop in X/A at $*$.
Case 1: Assume $a_0 = lim_{t\to 0}\widetilde{\alpha}(t)$ and $a_1 = lim_{t\to 1}\widetilde{\alpha}(t)$ exist, so a_0,$a_1 \in A$ and we can define a path $\overline{\alpha} : I \longrightarrow X$ such that $\overline{\alpha}|_{(0,1)} = \widetilde{\alpha}$, $\overline{\alpha}(0) = a_0$, $\overline{\alpha}(1) = a_1$. Since A is path connected, there exist paths $\lambda_0, \lambda_1 : I \longrightarrow A$ such that λ_0 is a path from a to a_0 and λ_1 is a path from a_1 to a. Therefore $\lambda_0 * \overline{\alpha} * \lambda_1$ is a loop at a such that $p_*([\lambda_0 * \overline{\alpha} * \lambda_1]) = [\alpha]$.
Case 2: If at least one of the above limits does not exist, then we make a sequence $\{[\alpha_n]\}_{n\in\mathbb{N}}$ in $Im(p_*)$ so that converges to $[\alpha]$. Without lost of generality, assume $a_0 = lim_{t\to 0}\widetilde{\alpha}(t)$ exists and $a_1 \in A$ is a limit point of $\widetilde{\alpha}((1/2,1))$ by Lemma 3.10. We can define a continuous map $\overline{\alpha} : [0,1) \longrightarrow X$ such that $\overline{\alpha}|_{(0,1)} = \widetilde{\alpha}$, $\overline{\alpha}(0) = a_0$. By hypothesis, there is a countable local base $\{O_i\}_{i\in\mathbb{N}}$ for a_1. Let $\{G_i\}_{i\in\mathbb{N}}$ be a sequence of open neighborhood of a_1 such that $G_i = O_1 \cap ... \cap O_i$. Since X is locally path connected and the G_i's are open neighborhoods of a_1, there exist path connected open neighborhoods $G'_i \subseteq G_i$ of a_1. Since the point a_1 is a limit point, there are $t_i \in (1/2,1)$ such that $\widetilde{\alpha}(t_i) \in G'_i$, $t_i < t_{i+1}$, $t_n \longrightarrow 1$ and there are paths $\gamma_i : [t_i,1] \longrightarrow G'_i$ from $\overline{\alpha}(t_i)$ to a_1, for all $i \in \mathbb{N}$. Since A is path connected, there exist paths $\lambda_0, \lambda_1 : I \longrightarrow A$ such that λ_0 is a path from a to a_0 and λ_1 is a path from a_1 to a. Let $\overline{\alpha}_n := \lambda_0 * \overline{\alpha}|_{[0,t_n]} \circ \xi_n * \gamma_n \circ \zeta_n * \lambda_1$, where $\xi_n : [0,1] \longrightarrow [0,t_n]$

and $\zeta_n : [0,1] \longrightarrow [t_n, 1]$ are increasing linear homeomorphisms. Note that every $\overline{\alpha}_n$ is a loop in X at a and if $\beta_n := p \circ (\gamma_n \circ \zeta_n)$, then

$$\alpha'_n := p \circ \overline{\alpha}_n = e_* * \alpha|_{[0,t_n]} \circ \xi_n * \beta_n * e_*$$

is a loop in X at $*$ and $p_*([\overline{\alpha}_n]) = [\alpha'_n]$. Define $\alpha_n : I \longrightarrow X/A$ by

$$\alpha_n(t) = \begin{cases} \alpha(t) & 0 \le t \le t_n \\ p \circ \gamma_n(t) & t_n \le t \le 1 \end{cases}$$

which is a loop at $*$ and $[\alpha'_n] = [\alpha_n]$. Thus it suffices to prove that $\alpha_n \longrightarrow \alpha$. If $\alpha \in \langle K, U\rangle$, where K is a compact subset of $[0,1]$ and U is an open subset of X/A, then
i) If $* \notin U$, then $K \cap \alpha^{-1}(*) = \varnothing$. Let $m \in \mathbb{N}$ such that $t_m \ge max\ K$. Since for each $t \in K$, $t \le t_m$, $\alpha_n(t) = \alpha(t) \in U$, for all $n > m$ which implies that $\alpha_n(K) = \alpha(K) \subseteq U$, for each $n > m$.
ii) If $* \in U$, then $p^{-1}(U)$ is an open neighborhood of A, thus there exists $m \in \mathbb{N}$ such that $G'_n \subseteq p^{-1}(U)$, for each $n \ge m$. Therefore $Im(\gamma_n) \subseteq G'_n$ which implies that $Im(p \circ \lambda_n) \subseteq U$. Thus for all $t \in K$ and $n \ge m$ we have

$$\alpha_n(t) = \begin{cases} \alpha(t) \in U & t \in [0, t_n] \\ (p \circ \gamma_n)(t) \in U & t \in [t_n, 1]. \end{cases}$$

Therefore for a semi-simple loop α in X/A we have $[\alpha] \in \overline{Im(p_*)}$. Similarly, for every loop α such that $\alpha^{-1}(\{*\})$ is finite, $[\alpha] \in \overline{Im(p_*)}$ which implies that the homotopy class of every geometrically finite loop belongs to $\overline{Im(p_*)}$ by Lemma 3.9.

Step Two: If α is not geometrically finite, W_α is countable since every open subset of I is a countable union of open intervals. Let $I_j = \overline{L_j}$ where $W_\alpha = \{L_j | j \in \mathbb{N}\}$ and let

$$\alpha_j(t) = \begin{cases} \alpha(t) & t \in I_1 \cup ... \cup I_j \\ * & otherwise, \end{cases}$$

then $[\alpha_j] \in \overline{Im(p_*)}$ since α_j's are geometrically finite.

Since $\overline{(\overline{Im(p_*)})} = \overline{Im(p_*)}$, it suffices to show that $\alpha_j \longrightarrow \alpha$. For, if $\alpha \in \langle K, U\rangle$ for a compact subset K of $[0,1]$ and an open subset U of X/A, then
i) If $* \in U$, then for each $t \in K$ and $j \in \mathbb{N}$, $\alpha_j(t)$ takes value $\alpha(t)$ or $*$ which in both cases belongs to U, so $\alpha_j(K) \subseteq U$, for all $j \in \mathbb{N}$.
ii) If $* \notin U$, then $K \cap \alpha^{-1}(\{*\}) = \varnothing$, so $K \subseteq \cup_j L_j$. By compactness of K we have $K \subseteq \cup_s L_{j_s}$, for $s = 1, 2, ..., n_K$. Let $M = max\{j_s | s = 1, 2, ..., n_K\}$, then $\alpha_j(K) = \alpha(K) \subseteq U$, for each $j \ge M$.

□

Corollary 3.12. *Let $A_1, A_2, ..., A_n$ be disjoint path connected, closed subsets of a first countable, connected, locally path connected space X. Then for every $a \in \bigcup_{i=1}^n A_i$,*

$$\overline{p_* \pi_1^{top}(X, a)} = \pi_1^{top}(X/(A_1, A_2, ..., A_n), *).$$

Proof. Let $p_i : X/(A_1, A_2, ..., A_{i-1}) \longrightarrow X/(A_1, A_2, ..., A_i)$. Since every point of each $p_{i-1}(A_i)$ has a countable local base in the connected, locally path connected space $X/(A_1, A_2, ..., A_{i-1})$, by Theorem 3.11 the result holds. □

Corollary 3.13. *Let $A_1, A_2, ..., A_n$ be disjoint path connected, closed subsets of a first countable, connected, locally path connected space X such that $X/(A_1, A_2, ..., A_n)$ is semi-locally simply connected, then for each $a \in \bigcup_{i=1}^n A_i$, $p_* : \pi_1(X, a) \longrightarrow \pi_1(X/(A_1, A_2, ..., A_n, *)$ is an epimorphism.*

Proof. Since $X/(A_1, A_2, ..., A_n)$ is connected, locally path connected and semi-locally simply connected space, $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is a discrete topological group which implies that $Im(p_*) = \pi_1(X/(A_1, A_2, ..., A_n), *)$ by Corollary 3.12. □

In the following example, we show that the condition locally path connectedness for X is necessary in Theorem 3.11.

Example 3.14. *Let* $X_1 = \{(x, sin(2\pi/x)) \in \mathbb{R}^2 | \ 0 < x \leq 1\}$, $X_2 = \{(x,y) \in \mathbb{R}^2 | \ x^2 + \frac{y^2}{4} = 1 \ , \ y \leq 0\}$, $X_3 = \{(x,0) \in \mathbb{R}^2 | \ -1 \leq x \leq 0\}$ *and* $A = \{(0,y) \in \mathbb{R}^2 | \ -1 \leq y \leq 1\}$. *If* $X = X_1 \cup X_2 \cup X_3 \cup A$, *then* $\pi_1(X, x_0) = 0$ *and* $\pi_1(X/A, *) \cong Z$. *Since* X/A *is locally path connected and semi-locally simply connected space,* $\pi_1^{top}(X/A, *)$ *is discrete which implies that* $\overline{p_*(\pi_1^{top}(X, x_0))} \neq \pi_1^{top}(X/A, *)$.

In the next example, we show that with conditions of Theorem 3.11, p_* is not necessarily an epimorphism and hence the hypothesis semi-locally simply connectedness in Corollary 3.13 is essential.

Example 3.15. *Let* $C_n = \{(x,y) \in \mathbb{R}^2 | \ (x - \frac{1}{n})^2 + y^2 = \frac{1}{n^2}\}$, *for* $n \in \mathbb{N}$, $HE_o = \bigcup_{n\in\mathbb{N}} C_{2n-1}$, $HE_e = \bigcup_{n\in\mathbb{N}} C_{2n}$ *and* $X = (HE_0 \times \{0\}) \cup (HE_e \times \{1\}) \cup A$, *where* $A = (\{(0,0)\} \times I)$. *One can easily see that* X/A *is the Hawaiian earing space. Let* α *be the loop in* X/A *that traverse* $p(C_1), p(C_2), ...$ *in ascending order. By structure of the fundamental group of the Hawaiian earing* [6], $[\alpha] \notin Im(p_*)$ *since if* $p_*([\beta]) = [\alpha]$, *the loop* β *must traverse infinitely many times* A *which is a contradiction to the continuity of* β.

Corollary 3.16. *Let* $A_1, A_2, ..., A_n$ *be subsets of a first countable, connected, locally path connected space* X *with disjoint path connected closure such that each* A_i *is open or closed. Then for* $a \in \bigcup_{i=1}^n A_i$,

$$\overline{p_*\pi_1^{top}(X, a)} = \pi_1^{top}(X/(A_1, A_2, ..., A_n), *).$$

Proof. If A is open and B is closed subset of X such that $\overline{A} \cap B = \varnothing$, then there exists a countable local base in $b \in B$ such that every basic subset of B does not intersect A. On the other hand, X/A is locally path connected. By the proof of Corollaries 3.5 and 3.12 the result holds. □

4. SOME APPLICATIONS

After that Biss [2] equipped the fundamental group of a topological space and named it topological fundamental group, Fabel [7] showed that topological fundamental groups can distinguish spaces with isomorphic fundamental groups. Hence studying the topology of fundamental groups seems important. In this section we specify the topology of $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ for some special cases.
By $(X, A_1, A_2, ..., A_n)$ we mean an $(n+1)$-*tuple* of spaces with one of the following conditions $(*)$:
(i): The A_i's are open subsets of X with path connected closure.
(ii): X is a connected, locally path connected, first countable space and the A_i's are open or closed subsets of X with disjoint path connected closure.

Theorem 4.1. *For an* $(n+1)$*-tuple of spaces* $(X, A_1, A_2, ..., A_n)$ *with assumption* $(*)$, *if* X *is simply connected, then* $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ *is an indiscrete topological group.*

Proof. Since X is simply connected, $\{[e_*]\}$ is a dense subset of $\pi_1^{top}(X/(A_1, A_2, ..., A_n))$, by Corollaries 3.5 and 3.16. For every $1 \neq [\alpha] \in \pi_1^{top}(X/(A_1, A_2, ..., A_n))$, the left multiplication $L_{[\alpha]}$ is a homeomorphism which implies that $\{[\alpha]\}$ is also dense. Hence every nonempty open subset of $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ contains every point of $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$. □

If $\mathcal{U}$ is an open cover of a connected and locally path connected space X, then the subgroup of $\pi_1(X, x)$ consisting of all homotopy classes of loops that can be represented by a product of the following type:

$$\prod_{j=1}^{n} u_j v_j u_j^{-1},$$

where the u_j's are arbitrary paths (starting at the base point x) and each v_j is a loop inside one of the neighborhoods $U_j \in \mathcal{U}$, is called the Spanier group with respect to $\mathcal{U}$, denoted by $\pi(\mathcal{U}, x)$ [13, 8]. Let $\mathcal{U}, \mathcal{V}$ be two open covers, and let $\mathcal{U}$ be a refinement of $\mathcal{V}$. Then $\pi(\mathcal{U}, x) \leq \pi(\mathcal{V}, x)$.This inclusion relation induces an inverse limit defined via the directed system of all covers with respect to refinement. H. Fischer et all [8] called such inverse limit the Spanier group of the based space (X, x) and denoted it by $\pi_1^{sp}(X, x)$. The authors [12] introduced Spanier spaces which are spaces such that thier Spanier groups is equal to their fundamental groups.

Also, the authors prove that for a connected and locally path connected space X, $\overline{\{1\}} \subseteq \pi_1^{sp}(X,x)$. Using this we have the following corollary that guaranty the existence of the Spanier spaces.

Theorem 4.2. *For an $(n+1)$-tuple of spaces $(X, A_1, A_2, ..., A_n)$ with assumption $(*)$, if X is simply connected, then $X/(A_1, A_2, ..., A_n)$ is an Spanier space.*

In the following example, we show that there exists a simply connected, locally path connected metric space X with a closed path connected subspace A such that X/A is not simply connected and by Theorem 4.1 $\pi_1^{top}(X/A, *)$ is an indiscrete topological group and hence is a Spanier space.

Example 4.3. *Using definitions of Example 3.15, let CHE_o and CHE_e be cones over HE_o and HE_e with height $\frac{1}{2}$ and let $X = CHE_o \cup CHE_e \cup A$. By the van Kampen theorem, X is simply connected, but X/A is not simply connected* [9].

Theorem 4.4. *For an $(n+1)$-tuple of spaces $(X, A_1, A_2, ..., A_n)$, if $\pi_1^{top}(X, a)$ is compact and $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is Hausdorff, then p_* is an epimorphism and so is a closed quotient map.*

Proof. $Im(p_*)$ is compact and therefore closed that since p_* is continuous. By Corollaries 3.5 and 3.16, $\pi_1^{top}(X/(A_1, A_2\ , ..., A_n), *) = \overline{Im(p_*)} = Im(p_*)$ which implies that p_* is onto. ☐

Theorem 4.5. *For an $(n+1)$-tuple of spaces $(X, A_1, A_2, ..., A_n)$, if $\pi_1^{top}(X, a)$ is compact and $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is Hausdorff, then either $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is a discrete topological group or uncountable.*

Proof. If $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ has at least one isolated point, then every singleton is open since left translations $L_{[\alpha]} : \pi_1^{top}(X/A, *) \longrightarrow \pi_1^{top}(X/A, *)$ by $L_{[\alpha]}([\beta]) = [\alpha*\beta]$, are homeomorphism for every $[\alpha] \in \pi_1^{top}(X/A, *)$. Thus $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is a discrete topological group. By Theorem 4.4, $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is compact and if has no isolated point, it is uncountable since a nonempty compact Hausdorff space without isolated points is uncountable [11, Theorem 27.7]. ☐

Corollary 4.6. *For an $(n+1)$-tuple of spaces $(X, A_1, A_2, ..., A_n)$, if $\pi_1^{top}(X, a)$ is a compact, countable quasitopological group, then either $X/(A_1, A_2, ..., A_n)$ is semi-locally simply connected or $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is not Hausdorff.*

Proof. Let $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ be Hausdorff, then p_* is onto by Theorem 4.4. Since $\pi_1^{top}(X, a)$ is countable, $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is countable. Theorem 4.5 implies that $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is a discrete topological groups. Hence $X/(A_1, A_2, ..., A_n)$ is semi-locally simply connected. ☐

Corollary 4.7. *For an $(n+1)$-tuple of spaces $(X, A_1, A_2, ..., A_n)$, if $\pi_1^{top}(X, a)$ is finite and $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is Hausdorff, then $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is finite and $X/(A_1, A_2, ..., A_n)$ is semi-locally simply connected.*

Proof. By Theorem 4.4, p_* is onto which implies that $\pi_1^{top}(X/(A_1, A_2, ..., A_n), *)$ is finite. The last assertion follows from Corollary 4.6. ☐

References

[1] A. Arhangelskii, M. Tkachenko, Topological Groups and Related Structures. *Atlantis Studies in Mathematics*, 2008.

[2] D. Biss, The topological fundamental group and generalized covering spaces, *Topology and its Applications.* 124 (2002) 355–371.

[3] J. Brazas, The topological fundamental group and free topological groups. *Topology and its Applications.* 158 (2011) 779–802.

[4] J.S. Calcut, J.D McCarthy, Discreteness and homogeneity of the topological fundamental group, *Topology Proc.* Vol. 34 (2009) 339–349.

[5] J.S. Calcut, R.E Gompf, J.D McCarthy, Quotient maps with the connected fibers and the fundamental group, arXiv:0904.4465v1.

[6] J.W. Cannon, G.R. Conner, The combinatorial structure of the Hawaiian earring group, *Topology and its Applications.* 106 (2000) 225–271.

[7] P. Fabel, Topological fundamental groups can distinguish spaces with isomorphic homotopy groups, *Topology Proc.* 30 (2006) 187-195.

[8] H. Fischer, D. Repovs, Z. Virk, A. Zastrow, On semilocally simply connected spaces, *Topology and its Applications.* 158 (2011) 397–408.

[9] H.B. Griffiths, The fundamental group of two space with a common point, *Quarterly Journal of Mathematics.* 5 (1954) 175–190.

[10] J. Morgan, I. Morrison, A van Kampen theorem for weak joins. *Proc. London Math. Soc.* 53 (1986) 562–576.

[11] J.R. Munkres, Topology: A First Course, second ed., *Prentice-Hall*, Upper Saddle River, NJ, 2000.

[12] A. Pakdaman, H. Torabi, B. Mashayekhy, Spanier spaces and covering theory of non-homotopically path Hausdorff spaces. Preprint.

[13] E.H. Spanier, Algebraic Topology, McGraw-Hill, New York, 1966.